



OPEN ACCESS

Original research

Occupation-specific risk estimates for suicide and non-fatal self-harm from a Swedish cohort of male construction workers followed 1987–2018

Kirsten Mehlig ¹, Kjell Torén ², Anthony D LaMontagne ³,
Viktoria Wahlström ⁴, Jenny Nyberg ^{5,6}, Margda Waern ^{7,8},
Maria Åberg ^{9,10}

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/oemed-2023-109246>).

For numbered affiliations see end of article.

Correspondence to

Dr Kirsten Mehlig, School of Public Health and Community Medicine, Institute of Medicine, University of Gothenburg, Gothenburg, Sweden; kirsten.mehlig@gu.se

Received 6 October 2023
Accepted 5 February 2024
Published Online First
28 February 2024

ABSTRACT

Objectives While suicidal behaviour has become less prevalent in non-manual workers in recent decades, rates have increased in manual workers. We aimed to identify occupations within the construction industry with excess risk of suicide and non-fatal self-harm.

Methods This cohort of Swedish construction workers comprises 389 132 individuals examined 1971–1993 and followed 1987–2018 using national hospital and cause of death registers. More than 200 job titles were merged into 22 occupational groups. For 296 891 men alive in 1987 and active in the construction sector, survival was calculated from baseline to first event of non-fatal self-harm or suicide and censored for emigration, long-term unemployment, disability pension, retirement, death from other causes or end of follow-up. HRs with 95% CIs were obtained from multiple Cox proportional hazard regression.

Results Overall, 1618 cases of suicide and 4774 events of non-fatal self-harm were registered. Self-harm before baseline was the single largest risk factor for suicide, HR 9.3 (95% CI 7.5 to 11.6). Compared with the overall mean, labourers and rock workers had excess risk for suicide, HR 1.4 (95% CI 1.1 to 1.7) and 1.5 (95% CI 1.0 to 2.3), respectively, while electricians, clerks and foremen had reduced risk. Labourers, concrete workers, sheet metal workers, painters, glaziers and the group 'other construction workers' were at increased risk for non-fatal self-harm. Almost all categories of manual workers were at increased risk for suicidal behaviour relative to clerks and foremen.

Conclusions Specific occupations within the construction sector were associated with excess risk for suicidal behaviour. Future studies should identify underlying risk factors to inform tailored interventions.

INTRODUCTION

Suicide rates vary across countries and time periods, with presently highest rates in European countries and lowest rates reported in the Eastern Mediterranean,^{1,2} and temporal maxima during, for example, economic crises.³ There are also pronounced sex differences, as suicide is more common among men and non-fatal self-harm among women.^{1,2,4} Men tend to under-report mental ill health, more seldom seek professional help and use more lethal suicide methods than women.⁵ Psychiatric disorder and

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Construction workers have excess risk for suicidal behaviour compared with other employees but information is lacking which specific occupations are at greatest risk within the sector.

WHAT THIS STUDY ADDS

⇒ Occupations with elevated risk of suicidal behaviour included labourers, rock workers, and sheet metal workers, with differential risks for suicide and non-fatal self-harm.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ The results show which occupations are at elevated risk for suicidal behaviour compared with the average risk for employees in the construction sector, and where prevention efforts should be intensified.

a history of suicide attempt are important predictors of suicidal behaviour.^{1,6,7} Recent reports from England and the USA show increasing trends in both suicide and non-fatal self-harm in the general population, indicating that non-fatal self-harm is both a risk factor and an important indicator for suicide risk.^{8,9} Large differences in suicide rates among employed men emphasise the importance of occupational aspects such as workplace conditions and job security.¹⁰ While early studies saw excess risk among men in professional occupations there was a reversal in trends, with increasing risk among manual workers in recent decades.^{11,12} A British study showed that suicide rates among manual workers were higher than among non-manual workers and professionals, and the rates among the former increased between 1979 and 2005, with labourers in building trades showing the second largest increase after coal miners.¹² An Australian study demonstrated that occupational differences in suicide risk were exacerbated during financial crises; among men, the lower skilled workers such as labourers were most affected.³ Low skill level is also associated with low job control and job insecurity, both of which are associated with suicide in general working populations.^{10,13–15} In Sweden, the



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY. Published by BMJ.

To cite: Mehlig K, Torén K, LaMontagne AD, *et al.* *Occup Environ Med* 2024;**81**:142–149.

age-adjusted suicide rate for men was 15.8 per 100 000 in 2016, which was higher than the global mean of 13.7 per 100 000 in men of all ages.¹⁶ Swedish conscription data from 1985 to 2005 showed that there was a growing social divide in physical health among the conscripts.¹⁷ Subsequent economic recessions may have further exacerbated social differences in general health, disproportionately affecting manual workers.³

Workers within the construction sector comprise a high-risk group regarding suicidal behaviour, as they are exposed to several risk factors such as heavy physical workload, strict deadlines, dangerous working conditions and toxic substances.^{1 18} Specific psychological risk factors in this male-dominated occupation are a low awareness of mental health problems and a certain bullying culture particularly directed towards apprentices and newcomers to the industry.^{19 20} Compared with the general working population, excess risk of suicide in construction workers has been demonstrated previously,^{11 12 21–23} but a detailed analysis of specific occupations within the sector is lacking. Also, occupational risk for non-fatal self-harm is an area not often researched in spite of its importance not least for subsequent suicide; reasons for this research gap may include lack of reliable outcome information, in particular as most people who self-harm do not seek medical attention,²⁴ which may be exacerbated among male construction workers for reasons given above. Episodes of self-harm among manual workers may also be misclassified as accidents, for instance, when heavy machinery is involved.²⁵

Thus, the aim of this study was to identify specific occupations within the construction industry where workers are at excess risk for suicide and non-fatal self-harm. Analyses were based on data from a historical cohort of Swedish construction workers followed up to 32 years or until the end of active work life. Valid outcome measures were obtained through linkage to nationwide hospital and cause of death registers.²⁶ Risk in individual occupations was compared with the average risk for suicidal behaviour in this sector, to describe the entire spectrum of risk estimates without singling out a specific reference occupation. The large size of almost 300 000 men allowed separate analyses of fatal and non-fatal self-harm in relation to 22 occupational groups.

METHODS

Study population and definition of analytical sample

Following an agreement between employers and unions, Swedish construction workers were affiliated with the National Occupational Health Service (Bygghälsan) that offered free health examinations to employees in the construction sector on a regular basis between the mid-1960s and 1993. Although the programme was voluntary, about 80% of all workers participated in at least one examination.²⁷ From 1971 on, data from medical examinations were entered into an electronic database. The current prospective study was based on data from 389 132 employees in the construction sector who participated in health examinations between 1971 and 1993, and were followed until 2018 using various national registers. Participants were re-examined up to 12 times after their first examination, with an average of three examinations in total. Observations from women who took part in the health examinations were excluded from this study because they constituted a minority (5%), and most were employed in administration (83% of all women). As most construction workers retired at age 65 during the study period, men who were aged 65 or above at their first health examination were excluded from the study. Further exclusions were made for men with missing occupational information. Because information on suicide was not available from the National Cause of Death register before 1987, the cohort was restricted to men alive and less than 65 years old in 1987. Lastly, we excluded all men who emigrated before 1987. The final analytical sample consisted of 296 891 male construction workers aged 16–64 (online supplemental table S1). For all participants, we had information on the year of birth as well as the region where the baseline examination was performed (14 regions from all over Sweden). It is noted that data were not collected for research purposes but to monitor individuals' health, and in contrast to health parameters, other information such as occupation was not always noted. Because the lack of occupational information can be assumed to be missing at random, the listwise deletion of observations with missing exposure values is not likely to cause bias in the association analyses presented below.

Exposure

More than 200 job titles within the construction sector were registered between 1971 and 1993.²⁸ These were combined into 22 groups of comparable task and skill level as proposed by technical experts from industry and unions, and included 19 groups of manual workers, clerks, foremen, as well as other workers within the construction sector (table 1). Detailed descriptions of the 19 manual work groups have been published previously.²⁹ In this study, the term 'labourers' is used to denote ground preparatory workers. Examples of job titles in the 'other construction workers' category are scaffold builders, welders, and

Table 1 Cohort description at baseline† and at last health examination

Occupation	Participants	Age	Participants at last health visit
	N (%)	Mean (SD)	N (%)
Road construction worker	3652 (1.2)	36.1 (10.8)	3671 (1.2)
Rock worker	2622 (0.9)	38.9 (10.4)	2404 (0.8)
Labourer	10215 (3.4)	36.2 (11.6)	10287 (3.5)
Concrete worker	28052 (9.5)	37.5 (12.4)	27079 (9.1)
Carpenter	62 628 (21.1)	32.7 (11.2)	61 795 (20.8)
Bricklayer	8514 (2.9)	35.6 (12.2)	8347 (2.8)
Floor layer	5183 (1.8)	32.7 (10.5)	5144 (1.7)
Glazier	2506 (0.8)	32.2 (10.7)	2482 (0.8)
Insulation worker	2679 (0.9)	32.8 (11.0)	2534 (0.9)
Sheet metal worker	11 353 (3.8)	31.2 (10.4)	11 009 (3.7)
Roofer	1289 (0.4)	34.5 (10.0)	1331 (0.5)
Pipe fitter, plumber	22 131 (7.5)	34.0 (11.6)	21 296 (7.2)
Painter	21 088 (7.1)	32.6 (11.3)	21 013 (7.1)
Machine operator	9819 (3.3)	37.2 (10.3)	9566 (3.2)
Crane operator	2886 (1.0)	39.9 (9.4)	2891 (1.0)
Driver	3998 (1.4)	38.6 (10.8)	3837 (1.3)
Refrigeration mechanic	1296 (0.4)	32.5 (10.0)	1288 (0.4)
Reparation mechanic	2594 (0.9)	36.9 (11.1)	2497 (0.8)
Electrician	34 712 (11.7)	31.3 (10.4)	33 673 (11.3)
Other construction work*	17 724 (6.0)	32.6 (12.5)	18 922 (6.4)
Foreman	29 002 (9.8)	38.4 (10.1)	32 238 (10.9)
Clerk, employee	12 948 (4.4)	41.0 (9.8)	13 587 (4.6)
Total	296 891 (100)	37.2 (13.4)	296 891 (100)

*Scaffold builder, welder, blacksmith, etc.

†Most recent health examination ≤1987 or directly after.

blacksmiths. Baseline occupation was defined as the occupation reported at the last examination before 1987, or, for those who entered the cohort after 1986, the occupation reported at the first examination. To account for the effect of switching among occupational groups, we also registered the occupation reported at the last health visit.

Outcomes and censoring information

Information on suicide deaths was obtained from the National Cause of Death Register and the National Hospital Register provided data on non-fatal self-harm. Self-harm is defined here as any type of self-injurious behaviour, including both suicide attempts and non-suicidal self-injuries.¹ Death by suicide and first episode of non-fatal self-harm were the main endpoints in this study and included events both with (International Classification of Diseases (ICD)-8/9: E95; ICD-10: X6, X7, X80–X84, Y870) and without (ICD-8/9: E98; ICD-10: Y1 Y2 Y30–Y34 Y872) determined intent. As some cases registered as self-harm of undetermined intent may have been accidental, we then carried out separate analyses after exclusion of cases with undetermined intent. Regarding non-fatal self-harm, we also distinguished between episodes before baseline examination and incident events. Observations were censored in connection with emigration due to lack of prospective information. Observations were also censored at the first time of unemployment and disability pension as both phenomena are associated with suicide¹ and can be expected to vary across occupations. Information on emigration and disability pension was obtained from the longitudinal integrated database for Health Insurance and Labour Market Studies (LISA) covering the years 1990–2018. The number of days on unemployment benefit was available from 1992 onwards and long-term unemployment was defined

as 1 year or more. The LISA register includes detailed data on health and social insurance and is updated yearly (Statistics Sweden SCB).

Statistical methods

To assess how occupation reported at baseline was associated with self-harm before baseline we used logistic regression and reported results as ORs with 95% CI. The prospective associations between 22 occupational groups and risk of suicide or non-fatal self-harm were examined using Cox proportional hazard models. Baseline was defined as 1 January 1987, or the date of the first health examination after that date. Observations were censored at first emigration after baseline, long-term unemployment, receipt of disability pension, age 65, death from other causes or end of 2018, whatever occurred first. Survival from baseline until the date of suicide, first record of non-fatal self-harm or censoring was calculated. For endpoints (suicide and non-fatal self-harm) with determined intent survival was calculated until the event of interest or censoring irrespective of previous events lacking determined intent. Men with a record of self-harm before baseline were excluded from prospective analyses of non-fatal self-harm to reduce the risk of previous mental health problems influencing the choice of occupation (reverse causation). Models for suicide were adjusted for self-harm before baseline, to account for previous mental health problems. As suggested by Stack,³⁰ regression models were adjusted for demographic variables, that is, age (with quadratic and cubic terms), region and year of occupational report. The proportional hazard assumption was tested and confirmed graphically for all occupational groups. Because endpoints were generally rare, we used penalised likelihood estimation to reduce the small sample bias in maximum likelihood estimation.³¹ Results were given in terms of HRs and CIs. As there is no natural reference category, we used deviation from means coding to calculate the risk for individual occupations compared with the overall risk for suicide or self-harm in the sample. This is done by obtaining linear combinations of occupation-specific beta-values such that their sum is zero, and the mean risk is given by the intercept in the logistic regression model and the baseline hazard in the survival model, respectively.³² Furthermore, we examined the association between the occupation reported at the last health examination and suicidal behaviour, again excluding all men with episodes of self-harm before the last health visit from the analysis of non-fatal self-harm. This was done to assess whether associations with the most recent occupation differed from results for baseline occupational exposure. Effect modification by self-harm before baseline was examined by adding product terms for self-harm before baseline and occupations to the model for suicide, and the p value for an overall F-test comparing models with and without interactions terms was given. Statistical analyses were performed by using SAS V.9.4 and Matlab (R2016b; The MathWorks). The significance level was set at 0.05 (two-sided tests).

RESULTS

Cohort description

Table 1 shows the number and mean age of workers, overall and by occupational group. At baseline, the mean age of construction workers was 37.2 years with differences between occupational groups. Among the workers with at least two examinations (64%), the mean number of changes between occupational categories was 0.3, range=0–7, and 77% did not change category. The right column of table 1 shows the distribution of occupational groups reported at the last health examination before the

Table 2 Occupation at baseline and odds for self-harm prior to baseline (n=2072/296 891)

Baseline occupation	No of cases (%)	OR (95% CI)‡
Road construction worker	29 (0.8)	1.02 (0.72 to 1.45)
Rock worker	33 (1.3)	1.52 (1.09 to 2.12)*
Labourer	124 (1.2)	1.49 (1.24 to 1.79)***
Concrete worker	295 (1.1)	1.29 (1.14 to 1.47)***
Carpenter	386 (0.6)	0.80 (0.71 to 0.90)***
Bricklayer	78 (0.9)	1.16 (0.93 to 1.44)
Floor layer	57 (1.1)	1.43 (1.11 to 1.85)**
Glazier	29 (1.2)	1.46 (1.03 to 2.08)*
Insulation worker	38 (1.4)	1.87 (1.37 to 2.56)***
Sheet metal worker	90 (0.8)	1.05 (0.85 to 1.29)
Roofer	19 (1.5)	1.84 (1.19 to 2.83)**
Pipe fitter, plumber	170 (0.8)	0.97 (0.83 to 1.14)
Painter	161 (0.8)	0.96 (0.81 to 1.13)
Machine operator	63 (0.6)	0.82 (0.64 to 1.05)
Crane operator	37 (1.3)	1.46 (1.07 to 2.00)*
Driver	21 (0.5)	0.69 (0.46 to 1.04)
Refrigeration mechanic	10 (0.8)	0.98 (0.54 to 1.75)
Reparation mechanic	16 (0.6)	0.84 (0.53 to 1.34)
Electrician	142 (0.4)	0.52 (0.44 to 0.61)***
Other construction work†	162 (0.9)	1.39 (1.18 to 1.64)***
Foreman	78 (0.3)	0.31 (0.25 to 0.39)***
Clerk, employee	34 (0.3)	0.33 (0.24 to 0.46)***

*p<0.05, **p<0.01, ***p<0.001.

†Scaffold builder, welder, blacksmith, etc.

‡Logistic regression adjusted for age, year, and region (ref=overall mean).

health programme ended in 1993. The prevalence of manual occupations remained stable, but the proportion of foremen increased by 1.1%.

Associations with self-harm before baseline

Episodes of self-harm before baseline were recorded for 2072 participants (0.3%). Several occupations showed excess risk for self-harm prior to baseline, that is, insulation workers, roofers, rock workers, labourers, crane operators, glaziers, floor layers, other construction workers and concrete workers, in order of decreasing risk compared with mean risk (table 2). Carpenters, electricians, clerks and foremen were less likely to have a history of self-harm compared with overall risk. To limit the potential for confounding, men with self-harm before baseline were excluded from the prospective analysis of non-fatal self-harm. For the same reason, the analysis of suicide was adjusted for self-harm before baseline.

Risk for suicide and self-harm among construction workers

Overall, 2624 cases of suicide and 7187 cases of non-fatal self-harm were recorded between baseline and end of follow-up in 2018. Censoring at age 65, emigration after baseline, unemployment and disability pension reduced the numbers to 1618 suicides and 4774 incident cases of non-fatal self-harm. Mean age at suicide was 46.7 years, and the corresponding figure for first episode non-fatal self-harm in men with no self-harm before baseline was 46.4 years. Average follow-up was 20.8 years for suicide (11.5 years among cases of suicide), and 20.6

years for non-fatal self-harm excluding observations with self-harm before baseline (15.5 years among cases of non-fatal self-harm).

Table 3 presents the associations between occupational categories and suicide and incident non-fatal self-harm after baseline, overall and restricted to endpoints with determined intent. Episodes of self-harm before baseline were a strong risk factor for suicide, HR=9.3 (95% CI 7.5 to 11.6). The HR remained high after censoring for suicide with undetermined intent, HR=8.0 (95% CI 6.2 to 10.5). Labourers, rock workers and other construction workers showed elevated risk for suicide compared with the overall mean, and these associations were particularly pronounced when deaths of undetermined intent were censored. The occupation-specific associations were not confounded by self-harm before baseline (not shown) nor were they modified by the latter (p value for interaction between occupation and self-harm before baseline in relation to suicide=0.7). Figure 1A shows that the occupation-specific risk estimates for suicide were hardly affected on exclusion of workers with previous self-harm. Sheet metal workers, other construction workers, painters, labourers and concrete workers showed elevated risk for non-fatal self-harm with determined intent but only for sheet metal workers, painters and other construction workers. Non-manual workers (clerks, foremen) as well as electricians had lower risks for suicide and self-harm compared with the overall mean. Figure 1 shows the consistency of occupation-specific risk across endpoints.

Table 3 Occupation at baseline and risk for suicide and non-fatal self-harm

Occupation	Suicide (n=296 891)		Non-fatal self-harm (n=294 819)	
	All cases (n=1618)	With known intent (n=1309)	All cases (n=4774)	With known intent (n=2169)
Occupation	HR (95% CI)†‡	HR (95% CI)†‡	HR (95% CI)†	HR (95% CI)†
Road construction worker	1.20 (0.82 to 1.74)	1.14 (0.75 to 1.74)	1.00 (0.78 to 1.28)	1.20 (0.85 to 1.69)
Rock worker	1.53 (1.00 to 2.33)*	1.85 (1.20 to 2.86)**	1.29 (0.94 to 1.77)	1.15 (0.72 to 1.84)
Labourer	1.35 (1.07 to 1.69)*	1.31 (1.01 to 1.70)	1.17 (1.00 to 1.36)*	1.15 (0.92 to 1.44)
Concrete worker	1.10 (0.92 to 1.30)	1.00 (0.82 to 1.22)	1.15 (1.03 to 1.27)*	1.11 (0.95 to 1.29)
Carpenter	0.97 (0.86 to 1.10)	0.98 (0.85 to 1.13)	1.00 (0.92 to 1.08)	0.91 (0.81 to 1.02)
Bricklayer	1.21 (0.93 to 1.57)	1.07 (0.78 to 1.46)	0.97 (0.82 to 1.16)	1.20 (0.95 to 1.51)
Floor layer	0.81 (0.55 to 1.18)	0.85 (0.56 to 1.28)	1.11 (0.91 to 1.34)	1.08 (0.81 to 1.44)
Glazier	0.92 (0.56 to 1.52)	0.84 (0.47 to 1.50)	1.16 (0.89 to 1.51)	1.40 (0.98 to 2.00)
Insulation worker	1.13 (0.73 to 1.77)	1.18 (0.73 to 1.92)	1.09 (0.83 to 1.43)	1.19 (0.82 to 1.74)
Sheet metal worker	1.16 (0.93 to 1.44)	1.10 (0.85 to 1.41)	1.23 (1.08 to 1.39)**	1.26 (1.05 to 1.51)*
Roofer	1.63 (0.95 to 2.79)	1.72 (0.96 to 3.09)	1.23 (0.84 to 1.82)	1.14 (0.64 to 2.05)
Pipe fitter, plumber	0.92 (0.76 to 1.11)	0.91 (0.74 to 1.12)	1.07 (0.96 to 1.19)	1.11 (0.95 to 1.29)
Painter	1.11 (0.93 to 1.32)	1.14 (0.93 to 1.39)	1.13 (1.02 to 1.25)*	1.39 (1.21 to 1.60)***
Machine operator	0.87 (0.66 to 1.14)	0.85 (0.62 to 1.16)	0.87 (0.73 to 1.04)	0.77 (0.59 to 1.02)
Crane operator	1.01 (0.62 to 1.64)	0.88 (0.49 to 1.58)	1.18 (0.86 to 1.62)	1.20 (0.77 to 1.87)
Driver	0.66 (0.40 to 1.09)	0.71 (0.42 to 1.22)	1.06 (0.81 to 1.39)	1.07 (0.74 to 1.57)
Refrigeration mechanic	1.18 (0.64 to 2.19)	1.16 (0.58 to 2.31)	0.72 (0.46 to 1.13)	0.81 (0.44 to 1.49)
Reparation mechanic	1.00 (0.61 to 1.65)	0.90 (0.50 to 1.62)	0.92 (0.66 to 1.28)	0.85 (0.51 to 1.39)
Electrician	0.73 (0.62 to 0.87)***	0.78 (0.65 to 0.94)**	0.70 (0.64 to 0.78)***	0.55 (0.47 to 0.65)***
Other construction work§	1.18 (0.97 to 1.45)	1.35 (1.09 to 1.66)**	1.19 (1.07 to 1.33)**	1.24 (1.06 to 1.46)**
Foreman	0.53 (0.43 to 0.66)***	0.57 (0.45 to 0.72)***	0.60 (0.52 to 0.68)***	0.50 (0.41 to 0.61)***
Clerk, employee	0.61 (0.46 to 0.83)**	0.64 (0.46 to 0.89)**	0.65 (0.54 to 0.79)***	0.56 (0.41 to 0.75)***

*p<0.05, **p<0.01, ***p<0.001.

†Cox proportional hazard regression adjusted for age, year and region (ref=overall mean).

‡Further adjusted for self-harm before baseline

§Scaffold builder, welder, blacksmith, etc.




Figure 1 Risk for suicide and non-fatal self-harm for manual and non-manual occupations within the construction sector (HRs from prospective analyses of suicide and non-fatal self-harm, ORs from cross-sectional analyses of self-harm before baseline; ref = overall mean risk).

Analyses using occupation at the last health examination, that is, the most recent occupation known in relation to outcome hardly changed the associations reported for baseline exposure, primarily due to the small number of changes in occupation recorded during 1971–1993 (online supplemental table S2). Roofers showed >60% excess risk for suicide compared with the overall mean, a result that was statistically significant only when considering occupations at last health visit. Reduced suicide risk for drivers was observed with respect to occupation reported at last visit only. Similarly, 45% excess risk for glaziers with respect to non-fatal self-harm was significant only if reported at last health visit. An overview of occupation-specific associations with suicidal endpoints is given in figure 2.

DISCUSSION

This study aimed to identify specific occupations within the construction sector with excess risk of suicidal behaviour compared with the mean risk in the entire cohort. Rock workers, labourers and roofers showed excess risk for suicide, while the higher skilled and higher status occupations of electricians, clerks and foremen were at reduced risk. Adverse associations increased in magnitude for rock workers and the group ‘other construction workers’ when analyses were restricted to suicide with determined intent. With the exception of labourers and the ‘other construction worker’ group the pattern differed with respect to non-fatal self-harm where excess risk was displayed by sheet metal workers, painters, glaziers and concrete workers.

This study showed that workers in specific occupational groups were at increased risk for different types of suicidal behaviour. Below, we offer possible explanations for the observed associations, which support the plausibility of results and may guide future studies and interventions. Compared with the overall mean electricians were the only manual workers who showed significantly lower risk for all types of suicidal behaviour, which may be related to higher skill level, better pay and better control over work tasks compared with other manual workers. In contrast, several occupations showed excess risk for specific endpoints, namely labourers, rock workers, and roofers for suicide, and sheet metal workers, painters, glaziers, and concrete workers for self-harm. A risk factor common to labourers, rock workers and roofers is the heavy physical labour that may lead to poor physical health including disability and chronic pain, which also affect mental health.³³ Previous findings from the same cohort confirm that these workers were more likely to receive disability pension compared with other construction workers.^{27 28} Labourers and rock workers were also among the oldest in this cohort, and thus more likely to be affected by the deterioration of physical health. Road construction work is characterised by heavy physical labour as well, yet it was not associated with any suicidal behaviour. A reason may be that the work is very intense during the ice-free season and with long holidays and a possibility for recovery during wintertime.

While excess risk for suicide among labourers and roofers has been shown previously,¹² our study provides further




Figure 2 Summary of significant occupation-specific associations with suicidal behaviours (occupations with excess risk in red, those with reduced risk in green; ref=mean risk for suicidal behaviour). ^aOnly observed for occupations reported at last health examination.

occupation-specific results for non-fatal self-harm. A study of 16-year old children from the British ALSPAC cohort showed differential risk factors for self-harm with and without suicidal intent, that is, higher values of IQ and maternal education for non-suicidal self-harm, and, among others, lower values of IQ and socioeconomic position for self-harm with suicidal intent.⁴ Although populations and outcome measures differ we note that the occupations associated with excess risk for suicide in our study (labourers, rock workers, roofers) were previously classified as low-skilled occupations while occupations associated with self-harm (sheet metal workers, painters, glaziers, concrete workers) were at higher skill level.²⁸ Since a positive correlation between IQ and professional skill level is plausible, our findings may provide further evidence for different risk factors behind the two kinds of suicidal behaviour including personality traits beyond IQ. Last, occupation-specific associations were largely consistent across non-fatal endpoints except for insulation workers, crane operators, glaziers and floor layers, who showed associations with self-harm before baseline, but no prospective associations. It is possible that the exclusion of individuals with episodes of self-harm before baseline reduced the possibility to capture prospective associations with non-fatal self-harm.

Strengths and limitations

The large, homogeneous cohort of construction workers is an important strength of the study that speaks for similar socio-cultural attitudes towards suicide.¹ The prospective design with long follow-up is a strength of the study, but the lack of occupational information after 1993 is a limitation as more recent

occupations may show a stronger link with suicidal behaviour, simply due to temporal proximity. We showed that using occupational information from the last health examination hardly changed the results, mainly because most workers kept their occupation while employed in the construction sector, but it is not clear whether this holds after 1993. Second, the lack of information from the cause of death registry before 1987 is a limitation that forced us to exclude observations from 4548 men, who died before 1987. This exclusion may have biased our results towards the null, in particular as the men, who died before 1987 were more likely to belong to risk groups, that is, rock workers, concrete workers, or workers from among the group of ‘other construction workers’, compared with the survivors (not shown). Last, these analyses are essentially descriptive, looking for subgroups with excess risk rather than causal factors. The observed associations could be due to selection into occupations or to the nature of and exposure in those occupations, or a combination of both. Even if occupation-specific risks are plausible, the lack of information on substance use and other mental disorders apart from a history of non-fatal self-harm is a limitation of this study as these risk factors may differ across occupational groups.^{18 34} However, a Danish study reported few occupational differences in suicide risk among people who suffered from psychiatric illness,³⁵ a finding that was paralleled in our study regarding self-harm before baseline.

Conclusions and implications for policy and practice

In this study, we identified specific high-risk occupations within the construction sector that should be prioritised for suicide

prevention. Our results are expected to be relevant today and across countries as tasks and conditions for construction work hardly vary, partly due to persistent economic difficulties that hamper the improvement of unfavourable work environments. Further research is needed to understand risk and protective factors in specific groups, to tailor interventions accordingly.

Author affiliations

- ¹School of Public Health and Community Medicine, Institute of Medicine, University of Gothenburg, Gothenburg, Sweden
- ²Occupational and Environmental Medicine, School of Public Health and Community Medicine, Institute of Medicine, University of Gothenburg, Gothenburg, Sweden
- ³Institute for Health Transformation, Deakin University, Geelong, Victoria, Australia
- ⁴Department of Public Health and Clinical Medicine, Umeå Universitet, Umeå, Sweden
- ⁵Section for Clinical Neuroscience, Institute of Neuroscience and Physiology, University of Gothenburg, Gothenburg, Sweden
- ⁶Sahlgrenska University Hospital, Region Västra Götaland, Gothenburg, Sweden
- ⁷Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, University of Gothenburg, Gothenburg, Sweden
- ⁸Department of Psychotic Disorders, Sahlgrenska University Hospital, Region Västra Götaland, Gothenburg, Sweden
- ⁹General Practice/Family Medicine, School of Public Health and Community Medicine, Institute of Medicine, University of Gothenburg, Gothenburg, Sweden
- ¹⁰Region Västra Götaland, Regionhälsan, Gothenburg, Sweden

Contributors MÅ conceived and designed the study in collaboration with MÅ, JN, KT, MW and VW. KM analysed the data and wrote the first draft, and all authors revised it critically for important intellectual content. All authors approved the final version to be published. KM is responsible for the overall content as guarantor. No AI technology has been used for any aspect of the study.

Funding This work was supported by AFA Insurance (an organisation owned by Sweden's labour market parties; MÅ, dnr 200265) and the Swedish state under the agreement between the Swedish government and the county councils, the ALF agreement (MÅ, ALFGBG-813511; MW, ALFGBG-965525).

Disclaimer The funding sources had no role in the study design, the collection, analysis and interpretation of the data, the writing of the report or the decision to submit the paper for publication.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and the Swedish Ethical Review Authority and Confidentiality Clearance at Statistics Sweden approved the study (reference number: Dnr 2020-01721). Also, the steering committee for the Bygghälsan register at Umeå University approved the study (01/20, 200402). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. Data will be made available on reasonable request following a decision by the steering committee.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

- Kirsten Mehlig <http://orcid.org/0000-0002-2653-0734>
 Kjell Torén <http://orcid.org/0000-0001-8509-7603>
 Anthony D LaMontagne <http://orcid.org/0000-0002-5811-5906>
 Viktoria Wahlström <http://orcid.org/0000-0002-4656-7606>
 Jenny Nyberg <http://orcid.org/0000-0002-4336-3886>
 Margda Waern <http://orcid.org/0000-0002-8330-6915>

Maria Åberg <http://orcid.org/0000-0002-0323-1061>

REFERENCES

- 1 Fazel S, Runeson B. Suicide. *N Engl J Med* 2020;382:266–74.
- 2 Turecki G, Brent DA, Gunnell D, et al. Suicide and suicide risk. *Nat Rev Dis Primers* 2019;5:74.
- 3 Milner AJ, Niven H, LaMontagne AD. Occupational class differences in suicide: evidence of changes over time and during the global financial crisis in Australia. *BMC Psychiatry* 2015;15:223.
- 4 Mars B, Heron J, Crane C, et al. Differences in risk factors for self-harm with and without suicidal intent: findings from the ALSPAC cohort. *J Affect Disord* 2014;168:407–14.
- 5 Mergl R, Koblberger N, Heinrichs K, et al. What are reasons for the large gender differences in the lethality of suicidal acts? An Epidemiological analysis in four European countries. *PLoS One* 2015;10:e0129062.
- 6 Erlangsen A, Appadurai V, Wang Y, et al. Genetics of suicide attempts in individuals with and without mental disorders: a population-based genome-wide Association study. *Mol Psychiatry* 2020;25:2410–21.
- 7 Favril L, Yu R, Uyar A, et al. Risk factors for suicide in adults: systematic review and meta-analysis of psychological autopsy studies. *Evid Based Ment Health* 2022;25:148–55.
- 8 Geulayov G, Kapur N, Turnbull P, et al. Epidemiology and trends in non-fatal self-harm in three centres in England, 2000–2012: findings from the Multicentre study of self-harm in England. *BMJ Open* 2016;6:e010538.
- 9 Martínez-Alés G, Keyes KM. Fatal and non-fatal self-injury in the USA: critical review of current trends and innovations in prevention. *Curr Psychiatry Rep* 2019;21:104.
- 10 Blomqvist S, Virtanen M, LaMontagne AD, et al. Perceived job insecurity and risk of suicide and suicide attempts: a study of men and women in the Swedish working population. *Scand J Work Environ Health* 2022;48:293–301.
- 11 Milner A, Spittal MJ, Pirkis J, et al. Suicide by occupation: systematic review and meta-analysis. *Br J Psychiatry* 2013;203:409–16.
- 12 Roberts SE, Jaremin B, Lloyd K. High-risk occupations for suicide. *Psychol Med* 2013;43:1231–40.
- 13 Milner A, LaMontagne AD. Suicide in the employed population: A review of epidemiology, risk factors and prevention activities. . 2018Available: <https://gupeaubguse/handle/2077/58195>
- 14 Åberg M, Staats E, Robertson J, et al. Psychosocial job stressors and risk of suicidal behavior - an observational study among Swedish men. *Scand J Work Environ Health* 2022;48:435–45.
- 15 Almroth M, Hemmingsson T, Kjellberg K, et al. Job control, job demands and job strain and suicidal behaviour among three million workers in Sweden. *Occup Environ Med* 2022;79:681–9.
- 16 Suicide in the world - Global Health Estimates: World Health Organization, . 2019Available: <https://www.who.int/publications/i/item/suicide-in-the-world>
- 17 Lissner L, Mehlig K, Rosengren A, et al. A growing social divide in body mass index, strength, and fitness of Swedish male conscripts. *J Adolesc Health* 2019;65:232–8.
- 18 Roche AM, Chapman J, Duraisingam V, et al. Construction workers' alcohol use, knowledge, perceptions of risk and workplace norms. *Drug Alcohol Rev* 2020;39:941–9.
- 19 Ross V, Mathieu SL, Wardhani R, et al. Factors associated with workplace bullying and the mental health of construction industry apprentices: A mixed methods study. *Front Psychiatry* 2021;12:629262.
- 20 Conway PM, Erlangsen A, Grynderup MB, et al. Workplace bullying and risk of suicide and suicide attempts: A register-based prospective cohort study of 98 330 participants in Denmark. *Scand J Work Environ Health* 2022;48:425–34.
- 21 Windsor-Shellard B, Gunnell D. Occupation-specific suicide risk in England: 2011–2015. *Br J Psychiatry* 2019;2019:1–6.
- 22 Andersen K, Hawgood J, Klieve H, et al. Suicide in selected occupations in Queensland: evidence from the state suicide register. *Aust N Z J Psychiatry* 2010;44:243–9.
- 23 Peterson C, Sussell A, Li J, et al. Suicide rates by industry and occupation - national violent death reporting system, 32 States, 2016. *MMWR Morb Mortal Wkly Rep* 2020;69:57–62.
- 24 Geulayov G, Casey D, McDonald KC, et al. Incidence of suicide, hospital-presenting non-fatal self-harm, and community-occurring non-fatal self-harm in adolescents in England (the iceberg model of self-harm): a retrospective study. *Lancet Psychiatry* 2018;5:167–74.
- 25 Robertson S. *Understanding men and health: masculinities, identity, and well-being*. Maidenhead: Open University Press, 2007.
- 26 Ludvigsson JF, Almqvist C, Bonamy A-KE, et al. Registers of the Swedish total population and their use in medical research. *Eur J Epidemiol* 2016;31:125–36.
- 27 Stattin M, Järholm B. Occupation, work environment, and disability pension: a prospective study of construction workers. *Scand J Public Health* 2005;33:84–90.
- 28 Järholm B, Stattin M, Robroek SJW, et al. Heavy work and disability pension - a long term follow-up of Swedish construction workers. *Scand J Work Environ Health* 2014;40:335–42.

- 29 Jackson JA, Olsson D, Punnett L, *et al.* Occupational Biomechanical risk factors for surgically treated Ulnar nerve Entrapment in a prospective study of male construction workers. *Scand J Work Environ Health* 2019;45:63–72.
- 30 Stack S. Occupation and suicide. *Social Science Quarterly* 2001;82:384–96. 10.1111/0038-4941.00030 Available: <https://onlinelibrary.wiley.com/doi/10.1111/0038-4941.00030>
- 31 Firth D. Bias reduction of maximum-likelihood-estimates. *Biometrika* 1993;80:27–38.
- 32 Hosmer DW, Lemeshow S. Applied logistic regression. In: *Applied Logistic Regression*. John Wiley & Sons, INC, 13 September 2000.
- 33 Jacobsen HB, Caban-Martinez A, Onyebeke LC, *et al.* Construction workers struggle with a high prevalence of mental distress, and this is associated with their pain and injuries. *J Occup Environ Med* 2013;55:1197–204.
- 34 Roche AM, Lee NK, Battams S, *et al.* Alcohol use among workers in male-dominated Industries: A systematic review of risk factors. *Safety Science* 2015;78:124–41.
- 35 Agerbo E, Gunnell D, Bonde JP, *et al.* Suicide and occupation: the impact of socio-economic, demographic and psychiatric differences. *Psychol Med* 2007;37:1131–40.

SUPPLEMENT**Table S1:** Definition of analytic sample

N = 389 132	Exclusions
	Women (n = 19 418)
N = 369 714	
	Age at baseline examination > 64 years (n = 6843)
N = 362 871	
	Occupation not reported at any health examination (n = 18 436)
N = 344 435	
	Age > 64 years before 1987 (n = 40 963)
N = 303 472	
	Death before 1987 (n = 4548)
N = 298 924	
	Emigration between last health examination and 1987 (n = 2033)
N = 296 891	

Table S2: Occupation registered at last health examination and risk for incident suicidal behavior

Occupation at last health visit	Suicide (n = 296 352)		Non-fatal self-harm (n = 294 156)	
	All cases (n = 1618)	With known intent (n = 1309)	All cases (n = 4647)	With known intent (n = 2086)
	HR (95% CI) ^{ab}	HR (95% CI) ^{ab}	HR (95% CI) ^a	HR (95% CI) ^a
Road construction worker	1.31 (0.91, 1.90)	1.19 (0.78, 1.82)	1.02 (0.80, 1.31)	1.28 (0.90, 1.81)
Rock worker	1.48 (0.95, 2.30)	1.77 (1.12, 2.80)*	1.28 (0.92, 1.78)	1.14 (0.69, 1.88)
Laborer	1.43 (1.15, 1.80)**	1.40 (1.09, 1.80)**	1.08 (0.92, 1.27)	1.07 (0.85, 1.35)
Concrete worker	1.10 (0.93, 1.31)	1.00 (0.82, 1.22)	1.19 (1.07, 1.32)**	1.23 (1.05, 1.43)**
Carpenter	0.99 (0.87, 1.13)	0.99 (0.86, 1.14)	1.00 (0.93, 1.08)	0.94 (0.84, 1.06)
Bricklayer	1.27 (0.98, 1.65)	1.13 (0.83, 1.54)	0.89 (0.74, 1.07)	1.10 (0.86, 1.41)
Floor-layer	0.83 (0.57, 1.21)	0.86 (0.57, 1.30)	1.10 (0.90, 1.33)	1.08 (0.80, 1.45)
Glazier	0.95 (0.58, 1.56)	0.85 (0.47, 1.53)	1.15 (0.87, 1.51)	1.45 (1.02, 2.07)*
Insulation worker	1.07 (0.67, 1.71)	1.08 (0.64, 1.82)	0.97 (0.72, 1.31)	0.99 (0.65, 1.53)
Sheet-metal worker	1.18 (0.94, 1.47)	1.08 (0.84, 1.40)	1.23 (1.08, 1.40)**	1.25 (1.03, 1.51)*
Roofer	1.69 (1.01, 2.84)*	1.63 (0.91, 2.94)	1.41 (0.99, 2.01)	1.25 (0.71, 2.18)
Pipe fitter, plumber	0.98 (0.81, 1.18)	0.97 (0.79, 1.19)	1.08 (0.97, 1.21)	1.15 (0.98, 1.34)
Painter	1.16 (0.97, 1.38)	1.18 (0.97, 1.43)	1.14 (1.03, 1.27)*	1.41 (1.22, 1.62)***
Machine operator	0.90 (0.69, 1.18)	0.86 (0.63, 1.17)	0.90 (0.76, 1.08)	0.85 (0.65, 1.11)
Crane operator	1.15 (0.73, 1.82)	1.12 (0.67, 1.88)	1.11 (0.80, 1.53)	1.12 (0.70, 1.79)
Driver	0.43 (0.23, 0.80)**	0.48 (0.25, 0.93)*	1.06 (0.81, 1.39)	1.03 (0.69, 1.52)
Refrigeration mechanic	1.49 (0.85, 2.60)	1.50 (0.81, 2.78)	0.88 (0.58, 1.33)	0.86 (0.47, 1.60)
Reparation mechanic	0.81 (0.46, 1.41)	0.83 (0.45, 1.53)	0.86 (0.61, 1.22)	0.82 (0.49, 1.37)
Electrician	0.77 (0.65, 0.91)**	0.81 (0.68, 0.97)	0.71 (0.64, 0.78)***	0.57 (0.49, 0.67)***
Other construction work ^c	1.16 (0.96, 1.41)	1.26 (1.02, 1.55)*	1.13 (1.01, 1.26)*	1.20 (1.01, 1.41)*
Foreman	0.57 (0.47, 0.70)***	0.61 (0.49, 0.75)***	0.64 (0.57, 0.72)***	0.49 (0.40, 0.60)***
Clerk, employee	0.50 (0.37, 0.68)***	0.53 (0.38, 0.74)***	0.63 (0.52, 0.76)***	0.56 (0.42, 0.75)***

^a Cox proportional hazard regression adjusted for age, year, and region (ref=overall mean)

^b Further adjusted for self-harm before baseline

^c Scaffold builder, welder, blacksmith etc.

* p <.05, ** p <.01, *** p <.001